




33. (Amended twice) A mixture comprising a set of at least six chemical compounds having a common heterocyclic scaffold bearing functionalizable atoms, wherein said set of compounds is represented by one of structures I, II or III:



III

wherein for structure I:

each tether moiety T is -NH(R¹)NH-, -NH(R¹)O-, -NHR²NH-, -NHR²SO₂NH-, -NHR¹-, -N(R⁴)₂, -N=N-, O, S, Se, -P(=O)(O)₂, NH, OR², OR³, malonato, pyrrolidinyl, piperidinyl, piperidinylmethylene, piperazinyl, or morpholino;

wherein R¹ is alkylene; R² is aryl; R³ is H or C₁-C₁₀ alkyl; R⁴ is alkyleneoxy; and

each chemical substituent L is, independently, C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, substituted C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, substituted C₂-C₁₀ alkynyl, C₄-C₇ carbocyclic alkyl, substituted C₄-C₇ carbocyclic alkyl, C₄-C₁₀ alkenyl carbocyclic, substituted C₄-C₁₀ alkenyl carbocyclic, C₄-C₁₀ alkynyl carbocyclic, substituted C₄-C₁₀ alkynyl carbocyclic, a nitrogen, oxygen or sulfur containing saturated heterocycle, a substituted nitrogen, oxygen or sulfur containing saturated heterocycle, a benzo-fused heterocycle, a substituted benzo-fused heterocycle, a substituted or unsubstituted saturated mixed

heterocycle; wherein each of the substituent groups is selected from a group consisting of alkyl, alkenyl, alkynyl, aryl, hydroxyl, alkoxy, benzyl, nitro, thiol, thioalkyl, thioalkoxy and halo; or

each chemical substituent L is, independently, substituted C_1 - C_{10} alkyl, wherein each of the substituent groups is selected from a group consisting of alkyl, alkenyl, alkynyl, aryl, hydroxyl, alkoxy, benzyl, nitro, thiol, thioalkyl, and thioalkoxy; or

L is, independently, piperazine, pyridazine, pyrazine, triazine, phthalimido, an ether having 2 to 10 carbon atoms and 1 to 4 oxygen or sulfur atoms, hydrogen, halogen, hydroxyl, thiol, keto, carboxyl, NR^1R^2 , $CONR^1$, amidine, guanidine, glutamyl, nitro, nitrate, nitrile, trifluoromethyl, trifluoromethoxy, NH-alkyl, N-dialkyl, O-aralkyl, S-aralkyl, NH-aralkyl, azido, hydrazino, hydroxylamino, sulfoxide, sulfone, sulfide, disulfide, silyl, a nucleosidic base, an amino acid side chain, or a carbohydrate; and

wherein for structures II and III:

each tether moiety T is $-NH(R^1)NH-$, $-NH(R^1)O-$, $-NHR^2NH-$, $-NHR^2SO_2NH-$, $-NHR^1-$, $-N(R^4)_2$, $-N=N-$, O, S, Se, $-P(=O)(O)_2$, NH, OR^2 , OR^3 , malonato, pyrrolidinyl, piperidinyl, piperazinyl, morpholino, imidazolyl, pyrrolyl, pyrazolyl, indolyl, 1H-indolyl, α -carbolinyl, carbazolyl, phenothiazinyl, phenoxazinyl, tetrazolyl, or triazolyl;

wherein R^1 is alkylene; R^2 is aryl; R^3 is H or C_1 - C_{10} alkyl; R^4 is alkyleneoxy; and

each chemical substituent L is, independently, C_1 - C_{10} alkyl, substituted C_1 - C_{10} alkyl, C_2 - C_{10} alkenyl, substituted C_2 - C_{10} alkenyl, C_2 - C_{10} alkynyl, substituted C_2 - C_{10} alkynyl, C_4 - C_7 carbocyclic alkyl, substituted C_4 - C_7 carbocyclic alkyl, C_4 - C_{10} alkenyl carbocyclic, substituted C_4 - C_{10} alkenyl carbocyclic, C_4 - C_{10} alkynyl carbocyclic, substituted C_4 - C_{10} alkynyl carbocyclic, C_6 - C_{14} aryl, substituted C_6 - C_{14} aryl, heteroaryl, substituted heteroaryl, a nitrogen, oxygen or sulfur containing heterocycle, a substituted nitrogen, oxygen or sulfur containing heterocycle, a mixed heterocycle, or a substituted mixed heterocycle; wherein each of the substituent groups is selected from a group consisting of alkyl, alkenyl, alkynyl, aryl, hydroxyl, alkoxy, benzyl, nitro, thiol, thioalkyl, thioalkoxy and halo; or